



**COMMITTEE ON HERBAL MEDICINAL PRODUCTS
(HMPC)**

DRAFT

**COMMUNITY HERBAL MONOGRAPH ON
PRIMULA VERIS L., *PRIMULA ELATIOR* (L.) HILL, RADIX**

DISCUSSION IN WORKING PARTY ON COMMUNITY MONOGRAPHS AND COMMUNITY LIST (MLWP)	January 2007 March 2007
ADOPTION BY HMPC FOR RELEASE FOR CONSULTATION	8 March 2007
END OF CONSULTATION (DEADLINE FOR COMMENTS)	15 June 2007
REDISCUSSION IN WORKING PARTY ON COMMUNITY MONOGRAPHS AND COMMUNITY LIST (MLWP)	
ADOPTION BY HMPC	

Comments should be provided using this [template](#) to hmpc.secretariat@emea.europa.eu
Fax: +44 20 75 23 70 51

KEYWORDS	Herbal medicinal products; HMPC; Community herbal monographs; traditional use; <i>Primula veris</i> L.; <i>Primula elatior</i> (L.) Hill; Primulae radix; primula root.
-----------------	---

**COMMUNITY HERBAL MONOGRAPH ON
PRIMULA VERIS L., *PRIMULA ELATIOR* (L.) HILL, RADIX**

1. NAME OF THE MEDICINAL PRODUCT

To be specified for the individual finished product.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION^{1, 2}

<u>Well-established use</u>	<u>Traditional use</u>
	<p>With regard to the registration application of Article 16d(1) of Directive 2001/83/EC as amended</p> <p><i>Primula veris</i> L. and/or <i>Primula elatior</i> (L.) Hill, radix (primula root)</p> <p>i) Herbal substance Whole or cut, dried rhizome and root</p> <p>ii) Herbal preparations A) Dry extract (3-9:1; 40-50 % v/v ethanol) B) Liquid extract (1:1; 70 % v/v ethanol), C) Liquid extract (1: 2.5; 70% v/v ethanol) D) Syrup (containing 1.5% dry extract A) E) Tincture (1:5; 70 % v/v ethanol) F) Soft extract (1-4:1, 20-55% v/v ethanol) G) Comminuted herbal substance</p>

3. PHARMACEUTICAL FORM

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Herbal substance or comminuted herbal substance for tea preparation or other herbal preparations in liquid and solid dosage forms for oral use. The pharmaceutical form should be described by the European Pharmacopoeia full standard term.</p>

¹ The material complies with the Eur. Ph. monograph (ref. 01/2005:1364)

² The declaration of the active substance(s) for an individual finished product should be in accordance with relevant herbal quality guidance.

4. CLINICAL PARTICULARS

4.1. Therapeutic indications

<u>Well-established use</u>	<u>Traditional use</u> Traditional herbal medicinal product used as an expectorant in cough associated with cold. The product is a traditional herbal medicinal product for use in specified indication exclusively based upon long-standing use.
-----------------------------	---

4.2. Posology and method of administration

<u>Well-established use</u>	<u>Traditional use</u> Posology <i>Adolescents over 12 years of age, adults, elderly</i> Single dose Herbal substance for tea preparation: 0.2 to 0.5 g Herbal preparations: A) Dry extract (according to ÖAB with DER 3-3.5:1): 0.1 – 0.2 g B) Liquid extract: 0.5 g D) Syrup: 5 – 10 ml E) Tincture: 0.5 – 1 g G) Comminuted herbal substance for tea preparation: 0.2 to 0.5 g Preparations A (different DER to ÖAB), C, F equivalent to the herbal substance Recommended mean daily doses Herbal substance for tea preparation: 0.5 to 1.5 g Herbal preparations: A) Dry extract (according to ÖAB with DER 3-3.5:1): 0.3 – 0.6 g B) Liquid extract: 1.5 g D) Syrup: 15 – 30 ml E) Tincture: 1.5 – 3 g G) Comminuted herbal substance for tea preparation: 0.5 to 1.5 g Preparations A (different DER to ÖAB), C, F equivalent to the herbal substance Dosage frequency: May be taken every 2 to 3 hours (up to a maximum 3 times daily)
-----------------------------	--

Children between 4 and 12 years of age

Herbal preparations:

C) Liquid extract

4-12 years of age

<i>Single dose</i>	<i>Dosage frequency</i>	<i>Daily dose</i>
0.3 ml	3 times daily	0.9 ml

E) Tincture

4-12 years of age

<i>Single dose</i>	<i>Dosage frequency</i>	<i>Daily dose</i>
0.3 ml	3 times daily	0.9 ml

F) Soft extract

4-6 years of age

<i>Single dose</i>	<i>Dosage frequency</i>	<i>Daily dose</i>
0.35 ml	3 times daily	1.05 ml

6-12 years of age

<i>Single dose</i>	<i>Dosage frequency</i>	<i>Daily dose</i>
0.35ml	3 to 4 times daily	1.05 ml to 1.4 ml

The use in children under 4 years of age is not recommended (see 4.4 Special warnings and precautions for use).

Duration of use

Children between 4 and 12 years of age

No longer than 5 days.

Adolescents over 12 years of age, adults, elderly

Medical attention should be sought if after 1 week of treatment the symptoms do not improve.

If the symptoms persist during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted.

Method of administration

Oral use

Tea preparation: 0.2 to 0.5 g of herbal substance or comminuted herbal substance for decoction, infusion or macerate.

As an expectorant one cup of tea every 2 to 3 hours.

4.3. Contraindications

<u>Well-established use</u>	<u>Traditional use</u> Hypersensitivity to the active substance or to other Primula species. Children with a history of acute stenosing laryngo-tracheitis. Asthma.
-----------------------------	--

4.4. Special warnings and precautions for use

<u>Well-established use</u>	<u>Traditional use</u> The use in children under 4 years of age is not recommended because medical advice should be sought. Caution is recommended in patients with gastritis or gastric ulcer. If dyspnoea, fever or purulent sputum occurs, a doctor or a qualified health care practitioner should be consulted. For tinctures and extracts, the appropriate labelling for ethanol, taken from the 'Guideline on excipients in the label and package leaflet of medicinal products for human use', must be included.
-----------------------------	---

4.5. Interactions with other medicinal products and other forms of interaction

<u>Well-established use</u>	<u>Traditional use</u> None reported.
-----------------------------	--

4.6. Pregnancy and lactation

<u>Well-established use</u>	<u>Traditional use</u> Safety during pregnancy and lactation has not been established. No adverse effects have been reported from the use of Primula root as a medicinal product during pregnancy and lactation. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.
-----------------------------	--

4.7. Effects on ability to drive and use machines

<u>Well-established use</u>	<u>Traditional use</u> No studies on the effect on the ability to drive and use machines have been performed.
-----------------------------	--

4.8. Undesirable effects

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Gastric disorders, nausea, vomiting and allergic reactions may occur. The frequency is not known.</p> <p>If other adverse reactions not mentioned above occur, a doctor or a qualified health care practitioner should be consulted.</p>

4.9. Overdose

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Overdose may lead to stomach upset, vomiting or diarrhoea.</p>

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC as amended.</p>

5.2. Pharmacokinetic properties

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC as amended.</p>

5.3. Preclinical safety data

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC as amended, unless necessary for the safe use of the product.</p> <p>Tests on reproductive toxicity, genotoxicity and carcinogenicity have not been performed.</p>

6. PHARMACEUTICAL PARTICULARS

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Not applicable.</p>

7. DATE OF COMPILATION/LAST REVISION

8 March 2007